

# EQUID HERPESVIRUS 1 & 4 MULTIPLEX qPCR TEST KIT USER GUIDE

# CAT NO.: YSL-qPX-IC-EHV-100

100 reactions with Internal Control and Lyophilised MasterMix

### **VERSION 5.5**

For research use only



YouSeq Ltd

8 Moorside Place Moorside Road Winchester SO23 7FX United Kingdom +44 333 577 6697 hello@youseq.com

youseq.com

Page 1



### INTENDED USE

This product is a qPCR test kit for detection of Equid herpesvirus 1 (EHV1) DNA and Equid Herpesvirus 4 (EHV4) DNA in good quality nucleic acid samples from a variety of sources. It is designed to be used by trained users in a suitable molecular biology laboratory environment.

### KIT CONTENTS

	Cap Colour	Volume
EHV1 specific primer/probe (FAM Probe) EHV4 specific primer/probe (ROX Probe) Internal extraction control primer/probe (VIC/HEX Probe)		110 µl
Lyophilised Tetra 2X qPCR MasterMix		1.1 ml*
EHV positive control template		500 μl*
Internal extraction control DNA template		500 μl*
MasterMix resuspension buffer		1.5 ml
DNase/RNase free water		1.5 ml
Template resuspension buffer		1.5 ml

 $^{*}$  Supplied lyophilised and requires resuspension before use, see resuspension step below for instructions

### RESUSPENSION

Resuspend the designated kit contents with the correct reagents as per the table below. Spin or gently tap the vials to ensure all the contents is at the bottom before opening.

After adding the resuspension reagent, pulse vortex the vials to ensure it is mixed well.

	Reagent	Volume
Lyophilised Tetra 2X qPCR MasterMix	MasterMix resus. buffer	1.1 ml
EHV positive control template	Template resus. buffer	500 µl
Internal extraction control DNA template	Template resus. buffer	500 µl





## MATERIALS REQUIRED BUT NOT PROVIDED

DNA Extraction kit - This qPCR test kit will work well with high quality DNA derived from any extraction kit with minimal PCR inhibitors present.

qPCR instrument with minimum 3 colour detection (FAM, ROX and VIC/HEX).

Pipettes, micro centrifuge tubes and general laboratory equipment.

### KIT SPECIFICITY

The YouSeq qPCR test kit for detection of Equid herpesvirus 1 & 4 (EHV) is designed to have the broadest detection profile possible and detect all clinically relevant strains. The primers and probes have very high (>95%) homology with all reference data within the NCBI database.

The target genes for Equid herpesvirus 1 and Equid Herpesvirus 4 (immediate-early protein (IE) gene and envelope glycoprotein C gene respectively) have been demonstrated to have unique sequences within these species making them ideal targets for highly specific detection of these pathogens.

If you require more specific data about the detection profile of the kit, please do not hesitate to contact our bioinformatics team : support@youseq.com

### USE OF DNA INTERNAL CONTROL

Add 5  $\mu$ l of the internal extraction control DNA into the extraction/lysis buffer for each sample that you are extracting. Make sure not to add this DNA directly in to your biological sample as this may cause degradation of the control DNA.

Continue DNA extraction as per the manufacturer's method.

### qPCR BENCH SIDE PROTOCOL

Clean and decontaminate all work surfaces, pipettes and other equipment prior to use to remove potentially contaminating nucleic acids.

### REACTION SET UP

Combine the following reagents to create a final test reaction:


Component	Volume
Tetra 2X qPCR MasterMix	10 µl
EHV specific primer/probe	1 μΙ
Extracted Sample DNA	9 µl
Final Volume	20 μl

Page 3



### **NEGATIVE CONTROL**

For a negative control reaction, repeat the reaction set up above replacing the sample DNA with DNase/RNase free water.



Please note: Make sure to seal the sample and negative control wells before proceeding to the positive control step.

### POSITIVE CONTROL

For a positive control reaction, repeat the reaction set up above replacing the sample DNA with 9  $\mu l$  of the positive control template supplied with the kit.

### qPCR AMPLIFICATION PROTOCOL

Run the following PCR protocol:

**Please note:** If using a qPCR machine that uses ROX as a passive reference, then the passive reference must be turned off or set to "none" indicating no passive reference.

1, , /	Temperature	Time
Hot Start	95°C	3 minutes
45 cycles	95°C	15 seconds
	60°C*	60 seconds

\*Make sure to collect fluorogenic data through FAM, ROX and VIC/HEX channels during this step

*
-
Daga 4

Page 4



### INTERPRETATION OF RESULTS

When analysing Sample Cq values, YouSeq recommends checking the threshold within the run file before interpreting the data. We would suggest setting the threshold to 10% of the relevant positive control End Point Fluorescence (EPF).

#### **Positive control**

Firstly, check the positive control performance. The positive control should amplify in a Cq range of approximately 18.5 + / -2. If the Cq range is not achieved, this would be a failed test and should be repeated.

Please note: The positive control in the kit is a representative sequence associated to the designs target region and does not contain the organisms entire genome.

The positive control does not include the internal control sequence. Therefore, the positive control should not be expected to amplify in the internal control channel.

#### **Negative control**

In ideal circumstances, the negative control should deliver a flat line – negative result. However, it is not uncommon for background laboratory contamination to cause a very late signal. If this signal is  $\geq$ 5 Cq values later than your sample signal then it can be considered negative and the result is viable.

If the negative control is <5 Cq later than the signal sample, then the result is inconclusive and the test should be repeated after potential sources of contamination have been removed.

The test is valid if the following conditions are met:



#### **Positive samples**

Samples that are positive for EHV1/EHV4 will deliver a defined "sigmoidal" amplification plot.

#### Internal extraction control

If your sample delivers a strong positive result then the internal extraction control is not required for data interpretation and can be ignored.

If your samples deliver a negative result, then the internal extraction control is useful to interpret the result. The Cq value from the internal extraction control will vary according to the amount of DNA in your sample. A late signal (Cq>28) indicates that only a small amount of host derived DNA was present in your sample. You may wish to repeat sample collection and then repeat the test in order to confirm the negative result.


	0
	•
and the second second	
	-1
	Page 5



# INTERPRETATION OF RESULTS CONTINUED

### Results interpretation at a glance:

	qPCR Signal				
EHV1 Sample (FAM Probe)	+	+	-	-	-
EHV4 Sample (ROX Probe)	+	-	+	-	-
Internal extraction Control (VIC/HEX Probe)	+/-	+/-	+/-	+	-
Result	Positive result EHV1 and EHV4 specific DNA detected	Positive result EHV1 specific DNA detected	Positive result EHV4 specific DNA detected	Negative Result	Failed test. Insufficient DNA



YouSeq Equid herpesvirus 1 & 4 Multiplex qPCR Test Kit with Internal Control and Lyophilised MasterMix Version 5.5

Page 6



# MULTIPLEX TROUBLESHOOTING

Тгасе	What can you see?	Cause	Action
2	One assay with greater end point fluorescence than another	Some fluorophores are brighter than others. Also, certain instruments detect different fluorophores with higher/lower efficiency	Analyse each channel individually so the Y- axis is correct for each fluorophore. or Analyse on logarithmic scale instead of linear scale
2 scence Units	Amplification/ unusual Cq value in unexpected fluorescent channel. Cq value/curve shape very similar to adjacent fluorescent channel	between channels. Amplification from one fluorescent channel has been mistakenly identified in its	Ensure manufacturer recommends the dye combination used in this kit. Recalibrate qPCR instrument

Page 7



### **PRODUCT SPECIFICATIONS**

#### Storing your kit

Store at -20°C from arrival. The qPCR kits shelf life is outlined as an expiry date on the pouch label.

#### **Use good quality DNA**

Poor quality input nucleic acid is the biggest cause of test failure. The kit will work well with any source of good quality DNA. Good quality is defined as DNA with high integrity (not degraded) and with low levels of inhibitors present.

#### **Regulatory status**

This product has been developed for Research Use Only and is not intended for diagnostic use. It should not be used for diagnosis of disease unless specifically approved by the regulatory authorities in the country of use.

#### **Quality Control**

In accordance with the YouSeq Ltd ISO EN 13485-certified Quality Management System, each lot of Equid herpesvirus 1 & 4 Multiplex qPCR kit is tested against predetermined specifications to ensure consistent product quality. Design of the kit met our robust bioinformatic analysis requirements resulting in a clinically relevant detection profile based on available sequence information. The kit is periodically checked against newly available sequence information to remain clinically relevant.

#### **Technical Assistance**

For customer support, please contact:

e-mail: support@youseq.com phone: +44 (0)333 577 6697

#### **Trademarks and Disclaimers**

#### YouSeq<sup>®</sup>, ROX<sup>™</sup>

Registered names, trademarks, etc. used in this document, even if not specifically marked as such, are not to be considered unprotected by law.

Not available in all countries. © 2023 YouSeq Ltd; all rights reserved.

Page 8